

Dirac's scalar field as dark energy within the frameworks of conformal theory of gravitation in Weyl–Cartan space

Olga V. Baburova*, Boris N. Frolov†, Roman S. Kostkin‡

*Moscow State Pedagogical University,
Department of physics for natural faculties,
MSPU, Krasnoprudnaya Street 14,
107140 Moscow, Russian Federation*

Abstract

Equations of the conformal theory of gravity with a Dirac scalar field in a Weyl–Cartan space-time have been derived. An exact solution of the equation for a scalar field, which has kind of a decreasing exponential function, has been found. This allows to explain a significant decrease (during of the period of inflation) the energy of physical vacuum (dark energy), which has been identified with the energy of the Dirac scalar field.

According to calculations of the quantum field theory, values of the cosmological constant responsible for accelerated expansion of the Universe should differ on 120 orders at early and modern stages of evolution. In [1], [2] one of the possible approaches has been developed to an explanation of such cosmological constant dynamics.

As the basis of the further constructions, we use Poincare–Weyl gauge theory of gravitation that has been developed in [3] – [5]. This theory is invariant both concerning the Poincare's subgroup and the Weyl subgroup – extensions and compressions (dilatations) of a spaces-time. In the theory there is the additional scalar field $\beta(x)$ introduced by Dirac. Its transformation under action of dilatations looks like: $\delta\beta = \varepsilon(x)\beta$.

*E-mail: baburova@orc.ru

†E-mail: frolovbn@orc.ru

‡E-mail: kostkin@orc.ru

As a consequence, an invariance of the theory under the following conformal transformations arises:

$$\begin{aligned}\delta g_{ab}^M &= 0, & \delta g_{\mu\nu} &= -2\varepsilon g_{\mu\nu}, & \delta h^a{}_\mu &= -\varepsilon h^a{}_\mu, & \delta \Gamma^a{}_{b\mu} &= \delta_b^a \partial_\mu \varepsilon, \\ \delta R^a{}_{b\mu\nu} &= 0, & \delta T^a{}_{\mu\nu} &= -\varepsilon T^a{}_{\mu\nu}, & \delta Q_\mu &= 8\partial_\mu \varepsilon.\end{aligned}$$

Here g_{ab}^M is the Minkowski metric of a tangent space, $T^a{}_{\mu\nu}$ is a torsion tensor, $Q_\mu = Q^\nu{}_{\nu\mu}$ is the Weyl's vector.

The Lagrange density of the theory in a Weyl–Cartan space-time is as follows:

$$\mathcal{L} = \mathcal{L}_G + \mathcal{L}_m + \frac{1}{2}\sqrt{-g}\Lambda^\mu{}_{ab}\left(Q^{ab}{}_\mu - \frac{1}{4}g^{ab}Q_\mu\right), \quad \Lambda^\mu{}_{ab}g^{ab} = 0,$$

where $\Lambda^\mu{}_{ab}$ are Lagrange multipliers, $\mathcal{L}_m$ is a Lagrange density of matter. The Lagrange density of a gravitational field we present as follows:

$$\mathcal{L}_G = \sqrt{-g}(f_0\beta^2 R + L_{R^2} + \beta^2 L_{T^2} + \beta^2 L_{Q^2} + \beta^2 L_{TQ} + L_\beta), \quad f_0 = \frac{1}{2\kappa}, \quad \kappa = 8\pi G, \quad c = 1. \quad (1)$$

It includes Lagrangians quadratic in curvature, L_{R^2} , and torsion, L_{T^2} , a Lagrangian L_{QT} containing a curvature–torsion interaction, and a proper Lagrangian of the scalar field

$$L_\beta = l_1 g^{\mu\nu} \partial_\mu \beta \partial_\nu \beta + l_2 \beta \partial_\mu \beta g^{\mu\sigma} T_\sigma + l_3 \beta \partial_\mu \beta g^{\mu\sigma} Q_\sigma + l_4 \beta \partial_\mu \beta Q^{\mu\sigma}{}_\sigma + \Lambda_0 \beta^4.$$

Variational equations of the field in the Weyl–Cartan space-time have been derived by variation of the full Lagrangian density of the theory. The independent variables are tetrads, a nonholonomic connection, the scalar field and the Lagrange multipliers.

At the first stage we neglect the contributions from terms quadratic in the curvature tensor, arising from the variation of the Lagrangian L_{R^2} .

Due to the cumbersome, the h –equation and the Γ –equation are not present here, while the β –equation has the form:

$$\begin{aligned}\frac{\delta \mathcal{L}}{\delta \beta} &= -2l_1 \overset{*}{\nabla}_\mu (\sqrt{-g} g^{\mu\nu} \partial_\nu \beta) - \beta \overset{*}{\nabla}_\mu (\sqrt{-g} (l_2 T^\mu + l_3 Q^\mu + l_4 Q^{\mu\lambda}{}_\lambda)) + \\ &+ 2\beta \sqrt{-g} \left(f_0 R + L_{T^2} + L_{Q^2} + L_{TQ} 2\Lambda_0 \beta^2 \right) + \frac{\delta \mathcal{L}_m}{\delta \beta} = 0,\end{aligned}$$

where $\overset{*}{\nabla}_\mu = \nabla_\mu + T_\mu$, $T_\mu = T^\nu{}_{\mu\nu}$. Variation of the Lagrange multipliers gives the Weyl condition on the nonmetricity tensor, $Q^{ab}{}_\mu = (1/4)g^{ab}Q_\mu$.

Variational equations are the equations of the gravitational field of the conformal theory of gravity in the Weyl–Cartan space-time with a Dirac scalar field in the tetrad

formalism. These equations are now investigated with a view to obtaining and solving an equation for the scalar Dirac field at the early stage of evolution of the universe in the absence of matter, $\mathcal{L}_m \equiv 0$.

In homogeneous and isotropic space-time the condition, $T^a_{\mu\nu} = -(2/3)h^a_{[\mu}T_{\nu]}$, takes place [6]. By means of this condition and in view of the Weyl condition, we can express from the Γ -equation a torsion trace and Weyl vector through the Dirac scalar field β :

$$T_\mu = \chi_T \partial_\mu \ln \beta, \quad Q_\mu = \chi_Q \partial_\mu \ln \beta. \quad (2)$$

The coefficients in (2) are expressed through couple constants of the original Lagrangian.

Considering the consequence of the h -equation together with the β -equation, and taking into account (2) and the condition of homogeneity and isotropy of space, we shall get,

$$\beta \ddot{\beta} - (k+1)(\dot{\beta})^2 = 0, \quad k = \frac{B}{A}, \quad (3)$$

where the constants A and B are expressed through the parameters of the original Lagrangian.

By replacing $\beta = u^{-1/k}$ (if $k \neq 0$) or $\beta = \exp(u)$ (if $k = 0$), this equation is reduced to the equation $\ddot{u} = 0$. Then the exact solution of equation (3) will be:

$$\beta = \frac{\beta_0}{(C_1 \hat{t} + 1)^{(1/k)}}, \quad (k \neq 0), \quad \beta = \beta_0 \exp(-C_2 \hat{t}), \quad (k = 0), \quad (4)$$

where β, C_1 and C_2 are arbitrary constants of integration. The constant β_0 is the value of the scalar field in the initial time, namely, in the Planck time. For reasons of quantum field theory, this quantity should be very huge. The constants C_1 or C_2 determine the initial value of the rate of the Dirac scalar field change.

Parameter k in rather complicated manner is determined by 16 of the coupling constants of the original Lagrangian (1). Therefore, we can always choose these constants so that one of the following conditions are fulfilled,

$$\frac{1}{k} = \frac{A}{B} \gg 1, \quad \text{or} \quad k = 0 \quad (B = 0). \quad (5)$$

Thus (at least for the open universe) if one of the conditions (5) is valid, one can provide the necessary rapid decrease in time the scalar field value.

The scaling-invariant theory of gravitation with scalar field in Riemann space-time has been developed in [7]–[9] (see also the references therein) in order to derive an alternative scenario of the evolution of the universe. In this scenario some of the observation data of modern cosmology can be explained without introducing Λ -term and without adopting

the inflation hypothesis. In contradiction with this, we do not reject inflation and Λ -term. In [1], [2], [10] we have expressed the hypothesis that the dark energy is determined by the value of the Dirac scalar field and determined by the term $\Lambda_0\beta^4$ of the Lagrangian.

Thus our result can explain the rapid decrease in time of the dark energy (the energy of physical vacuum). The decrease of dark energy according to (4) is much more intense than the corresponding decrease, which can be carried out in Poincare gauge theory of gravity [11], [12], in which the effective cosmological term is defined by the torsion tensor.

We point out that the ultra-rapid decrease of the energy of physical vacuum according to the law (4) occurs only prior to the Friedmann era evolution of the universe. According to the scenario of inflation, the birth of the rest masses of elementary particles begins in the last period of inflation, and further evolution of the universe is determined not by a scalar field, but mainly by the born ultrarelativistic matter and the radiation interacting with it. Solution (5) requires a suitable modification in this case.

References

- [1] Babourova O. V., Kostkin R. S., Frolov B. N. *Dirac scalar field in Weyl–Cartan space*. In: II Russian summer school-seminar - Modern theoretical problems of gravitation and cosmology (GRACOS-2009, August 24-29, Kazan-Yalchik). Proceedings of the school-seminar (OOO “Foliant”, Kazan, 2009). P. 127 (in Russian, <http://rgs.vniims.ru/main.htm>).
- [2] Babourova O. V., Kostkin R. S., Frolov B. N. *Cosmology of early universe in frameworks of conformal theory of gravitation in Weyl–Cartan space*. In: Matematika, Informatika, Fizika i ih Prepodavanie (MPGU, Moscow, 2009). P. 172–174 (in Russian).
- [3] Babourova O. V., Frolov B. N., Zhukovsky V. Ch. *Gauge Field Theory for Poincaré Weyl Group*. Phys. Rev. D . 2006. V. 74. P. 064012-1-12 (gr-qc/0508088, 2005).
- [4] Babourova O. V., Frolov B. N., Zhukovsky V. Ch. *A Weyl–Cartan space-time model based on the gauge principle*. Theor. Math. Phys. **157**(1), 1420–1432 (2008).
- [5] Babourova O. V., Frolov B. N., Zhukovsky V. Ch. *Theory of Gravitation on the Basis of the Poincaré Weyl Gauge Group*. Gravit. Cosmol. 2009. V. 15. No. 1. P. 13–15.
- [6] Tsamparlis M. *Cosmological principle and torsion*. Phys. Lett. 1979. V. 75A. P. 27–28.

- [7] Pervushin V., Proskurin D. *Conformal General Relativity*. Grav. Cosmol. Suppl. V. 8N1. 2002. P. 161–167.
- [8] Glinka L.A., Pervushin V.N. *Hamilton Unification of General Relativity and Standard Model*. The Old and New Concepts of Physics. 2008. V. 5. N1. P. 31–155.
- [9] Behnke D., Blaschke D., Pervushin V. N., Proskurin D. *Description of Supernova Data in Conformal Cosmology without Cosmological Constant*. Phys. Lett. V. B530. 2002. P. 20–26.
- [10] Frolov B. N., Kouchoumov A. Ju. *Weyl's nonmetricity and Dirac's scalar field as two components of dark energy. I. Effective cosmological term in theory of gravity with scalar field in weyl-cartan spacetime*. In: Physical Interpretations of Relativity Theory (Proc. XV Int. Sci. Meeting PIRT-2009) (Bauman Moscow State Technical University, Moscow, 2009). P. 359–366.
- [11] Minkevich A. V., Garkun A. S., Kudin V. I. *Regular accelerating Universe without dark energy*. Class. Quant. Grav. 2007. V. 24. P. 5835–5848 (gr-qc/0706.1157).
- [12] Minkevich A. V. *Accelerating universe with spacetime torsion but without dark matter and dark energy*. Phys. Lett. B. 2009. V. 678. P. 423–426 (gr-qc/0902.2860v2).